

Comment on "Liquid Limits: The Glass Transition and Liquid-Gas Spinodal Boundaries of Metastable Liquids"

In a recent paper¹, Sastry studies by numerical simulations the phase diagram of a simple fragile glass-forming liquid, the binary mixture with Lennard-Jones (LJ) interactions introduced by Kob and Andersen, presenting very interesting and clear results. Among other things, he obtains the position of the liquid-glass transition temperature as function of density ρ . He studies both the temperature $T_0(\rho)$ where the diffusion coefficient (of A particles) drops to zero according to the Vogel-Fulcher-Tammann-Hesse law, and the temperature where the extrapolation of data on the configurational entropy $S_c(\rho, T)$ vanishes, S_c being evaluated as the difference between the liquid entropy and the harmonic amorphous solid entropy. These two ways give values remarkably close.

We apply to the LJ binary mixture, at various density values, the analytic approach to structural glass thermodynamics recently introduced², which allows to obtain both the liquid-glass transition temperature T_K and properties in the glassy phase below T_K doing a first principles computation. In one version of this approach, which implements analitically the proposal of inherent structure dominance at low temperature, one needs to compute the liquid free-energy density f and two-point correlation functions $g(r)$ at $T \gtrsim T_K$.

The theory and different ways for implementing it are discussed in detail elsewhere²⁻⁵. We only recall that, within this framework, at T_K the system undergoes a second order transition corresponding to a one step replica symmetry breaking. As one could expect intuitively, it is found that in the region $T \gtrsim T_K$, where the phase space can be partitioned in 'valleys', the entropy of the liquid corresponds to the entropy of a typical valley S_{sol} plus the configurational entropy S_c , and the Kauzmann temperature is defined by the condition $S_c(T_K) = 0$.

We use the harmonic approximation scheme, which gives a simple formula for the single valley entropy in the liquid phase:

$$S_{sol} = \frac{3N}{2} \ln(2\pi e T) - \frac{1}{2} \langle \text{Tr} \ln \mathcal{M} \rangle,$$

where $\mathcal{M}$ is the $3N \times 3N$ Hessian (i.e. the matrix of second derivative of the potential with respect to the Nd particle coordinates) and $\langle \cdot \rangle$ means the Boltzmann average. We are therefore approximating S_{sol} by only keeping the harmonic vibrational contribution. Further approximations are needed for computing the spectrum. Here we consider fluctuations of the diagonal part of $\mathcal{M}$ up to second order⁵, while the whole nondiagonal contribution is summed up by means of a kind of chain approximation.

In the computation of the liquid free energy and of the correlation functions in the high density region we use the Hypernetted Chain Approximation (HNC). Unfortunately this approximation overestimates the gas-liquid transition temperature and a more refined one is neces-

sary at low densities. Liquid quantities at $\rho = 1.2$ have been computed⁵ within a scheme similar to that introduced by Zerah and Hansen (ZH), which also has the advantage of reducing the error in the approximation (of order 10% in the HNC case).




FIG. 1. The data on T_0 obtained by Sastry¹ compared with our analytical results.

We plot in [Fig. 1] the theoretical predictions on T_K at different density values, compared with the data on T_0 by Sastry. Despite of the error in liquid quantities obtained by HNC, there is a quantitative agreement which makes evident that the analytic approach works well and it can be used as a powerful tool for a better understanding of glassy systems.

To conclude, we stress that, at least in this case, the Kauzmann temperature turns out to be directly comparable with the temperature where the diffusion drops to zero, which strongly supports the theoretical expectation of an effective ergodicity breaking.

B. Coluzzi^(a), G. Parisi^(a) and P. Verrocchio^(b)

a) Dipartimento di Fisica, INFN and INFM, Università di Roma *La Sapienza*, P.le A. Moro 2, 00185 Roma, Italy.

b) Dipartimento di Fisica, Università di Trento, I-3805 Povo, Trento, Italy.

PACS numbers: 64.70.Pf, 63.50.+x

¹ S. Sastry, cond-mat/0005372 to appear on *Phys. Rev. Lett.*

² M. Mézard and G. Parisi, *Phys. Rev. Lett.* **82** 747 (1998); *J. Chem. Phys.* **111** 1076 (1999).

³ M. Mézard, *Physica A* **265**, 352 (1999); *First steps in glass theory* cond-mat/0005173.

⁴ B. Coluzzi, M. Mézard, G. Parisi and P. Verrocchio, *J. Chem. Phys.* **111** 9039 (1999).

⁵ B. Coluzzi, G. Parisi and P. Verrocchio *J. Chem. Phys.* **112** 2933 (2000); *Phys. Rev. Lett.* **84** 306 (2000).